

# Taking messages about breast cancer and smoking online:

New directions for reducing tobacco exposure among girls

Joan L. Bottorff, Jodie Stevens, Laura L. Struik and Chris G. Richardson



### Smoking affects more than lungs...

■ Tobacco smoke contains approximately 20 mammary carcinogens

(California Environmental Protection Agency, 2005).

- Exposure to tobacco smoke linked to premenopausal breast cancer (Collishaw et al., 2009; Johnson, 2012).
- Vulnerability high during breast cell growth (Innes & Byers, 2001; Lash & Aschengrau, 1999; Okasha, McCarron, Gunnell, & Smith, 2003).





#### **START**

**Goal:** To raise awareness among youth about girls' increased risk for breast cancer (BC) when exposed to tobacco smoke

#### **Objectives:**

- ☐ To develop gender- and Aboriginal-tailored messages
- ☐ To evaluate youths' responses to tailored messages





# **Program of Research**

#### **START** is nested in the British Columbia Adolescent Substance Use Survey (BASUS)

BASUS	Wave 2 (Spring 2010)	Wave 3 (Fall 2010)	Wave 4 (Spring 2011)	Wave 5 (Fall 2011)	Wave 6 (Spring 2012)
START	Phase 1 BASUS data used to identify and characterize groups based on smoking status, gender and ethnicity	Phase 2 Message development (Qualitative; focus groups)	using a ran		al (including a ted within the



### Message Development Phase

- Message strategies developed via 8 focus groups
  - 43 youth aged 12 to 17
    - 18 female
    - 20 of Aboriginal descent



**Be Breast Friends for Life** 







#### Girls' Messages

#### Smoking affects more than your lungs.

Cigarette smoke, even second hand smoke, puts girls at risk of breast cancer at an early age.



START START

If you smoke, think about quitting. Do it for yourself and for all the girls you know.

#### Smoking affects more than your lungs.

Cigarette smoke, even second hand smoke, puts girls at risk of breast cancer at an early age.



#### **Boys' Messages**





# **Standard Message**





#### Measures

- ☑ Demographics
- ☑ Smoking behaviour
- ✓ Message viewing time
- Perceived risk
- ✓ Worry
- ✓ Information-seeking






Participant Characteristics		Intervention n (%)	Control n (%)
Gender*	Male	310(48)	348(42)
	Female	339(52)	491(59)
Age in years***	13	92(14)	172(21)
	14	351(54)	480(57)
	15	206(32)	187(22)
Ethnicity	Aboriginal	71(11)	96(11.8%)
	Other	555(89)	717(88)
Family member has had BC*	Yes	153(25)	154(20)
Have tried smoking tobacco-	Yes	60(9)	104(12)
Does anyone smoke in your home?*	Yes	60(9)	107(13)
Past month's exposure to SHS	Every day	20(3)	35(4)
	Almost every day	70(11)	79(10)
	At least once a week	153(24)	236(28)
	At least once in past month	281(44)	347(43)
	Never	110(17)	119(15)
Tobacco identified as a risk factor for	Yes	172(27)	208(25)
BC			

# **Exposure to intervention**





# Time spent reading messages



#### Post-intervention relative risk

0.84 € My smoking increases my risk of BC (smoking girls) If I start smoking, it will increase my risk of BC (non-1.14 €€ \*\* smoking girls) 1.14 € \*\* Exposure to SHS increases my risk of BC (all girls) 1.14 €€ \*\* Exposure to SHS increases girls' risk of BC (all girls) Exposure to my SHS increases BC risk of the girls I 1.13€ spend time with (smoking girls) 1.10 €€ \* Exposure to SHS increases girls' risk of BC (all boys) Exposure to my SHS increases BC risk of the girls I 1.10<sup>€</sup> spend time with (smoking boys) 0.00 0.50 1.00 1.50

<sup>&</sup>lt;sup>€€</sup> Adjusted RR for age, family history of BC, smoking status of family and peers, and having tried cigarettes. \*p<0.01, \*\*p<0.001



<sup>€</sup> Unadjusted RR due to small sample size

## **Post-intervention worry**

I am worried that I could get BC because I am exposed to someone else's SHS (all girls)

I am worried that I could get BC because of my cigarette smoking (smoking girls)


I am worried that I could get BC if I started smoking cigarettes (non-smoking girls)

I am worried that SHS from other peoples' cigarette smoke increases girls' risk for getting BC (all girls)

I am worried that SHS from my ciagrette smoking increases the BC risk of the girls that I spend time with (smoking girls)

I am worried that SHS from other peoples' cigarette smoke increases girls' risk for getting BC (all boys)

I am worried that SHS from my cigarette smoking increases the BC risk of the girls that I spend time with (smoking boys)




\*p<0.01

0.00 0.20 0.40 0.60 0.80 1.00 1.20 1.40 1.60



### Interest in receiving more information





# Making sense of the results...

- Positively framed, gender-specific messages appear to increase awareness
- Girls' response may be related to physical/social changes with transition to womanhood
- Masculine norms can be repositioned to promote young men's involvement in girls/women's health issues
- Interactive technologies hold promise for cost-effective, gendered health promotion









# **Knowledge Translation**

- Information for girls and boys
  - About risk
  - Strategies to decrease girls' risk





# **Teaching Kits**

PowerPoint presentation and lesson plan for teachers



# Online resources for girls and boys





### **ePostcards**

Sent by email or though social media websites











# Girls' Feedback





# Boys' Feedback







- Interactive online quizzes
- Infographics
- You Tube video the biology behind the risk
- You Tube video breast cancer awareness for young women





Conclusion



# Acknowledgements

Additional team members
 R. Haines-Saah, J.L. Oliffe, P. Hutchinson, P.A. Ratner, C.T.C. Okoli, C. Gotay, K. Johnson, J. Schwartz, J. Memetovic



#### Funding for the project

- Canadian Breast Cancer Research Alliance
- Canadian Breast Cancer Foundation
- Career support for Chris Richardson: MSFHR, CIHR

#### Graphic design

Think Media, Media Button











Questions?

### For further information:

Dr. Joan L. Bottorff

**Institute for Healthy Living and Chronic Disease Prevention** 

**UBC's Okanagan Campus, Kelowna** 

E: Joan.bottorff@ubc.ca

**Dr. Chris Richardson** 

School of Population and Public Health UBC's Vancouver campus

E: Chris.richardson@ubc.ca





### References

- Bottorff, J.L., McKeown, S.B., Carey, J., Haines, R., Okoli, C., Johnson, K., Easley, J., Ferrence, R., Baillie, L., & Ptolemy, E. (2010). **Young women's responses to smoking and breast cancer risk information**. Health Education Research. 25(4), 668-677.
- Collishaw NE, Boyd NF, Cantor KP, Hammond SK, Johnson KC, Millar J, et al. (2009) Canadian Expert Panel on Tobacco Smoke and Breast Cancer Risk. [http://www.otru.org/pdf/special/expert panel tobacco breast cancer.pdf]
- Haines, R.J., Bottorff, J.L., Barclay McKeown, S., Ptolemy, E., Carey, J. & Sullivan, K. (2010). **Breast cancer messaging for younger women: gender, femininity and risk**. Qualitative Health Research, 20, 731-742.
- Johnson KC, Miller AB, Collishaw NE, Palmer JR, Hammond KS, Salmon AG, Cantor KP, Miller MD, Boyd NF, Millar J, Turcotte F: (2009) Active smoking and secondhand smoke increase breast cancer risk: the report of the Canadian Expert Panel on Tobacco Smoke and Breast Cancer Risk. Tobacco Control 2011, 20(e2).
- Johnson KC (2012). **Tobacco smoke and breast cancer risk: rapid evolution of evidence and understanding in the early 21st century.** In *Cigarette Consumption and Health Effects.* Edited by Chen GG. Ottawa, ON: Nova Science Publishers, Inc.(p.1-19).
- Struik, L., Bottorff, J.L., Jung, M., & Budgen, C. (2012). **Reaching girls through social networking: A venue for smoking prevention messages**. Canadian Journal of Nursing Research, 44(3), 84-103.

